

SURGICAL COMPOSITIONS AND METHODS OF USING THE SAME
DEXTRIN-CONTAINING COMPOSITION FOR PREVENTING SURGICAL ADHESIONS

PCT Rec'd 10 NOV 2000

This invention relates to the prevention of surgical adhesions, and in particular to adhesions taking place in serous cavities including the peritoneum, the pericardium, the plura and synovial cavities such as joints and tendons and to adhesions following spinal and/or cranial operations. Reference will be made hereinbelow to the prevention of adhesions in the peritoneum but it should be understood that the present invention has applicability in connection with other serous cavities in both humans and animals.

10

Abdominal surgery is a rapidly changing field. Many forms of open surgery are being increasingly replaced by laparoscopic procedures. Although considerable immediate post-surgical benefits have been demonstrated to follow from laparoscopic surgery, the incidence of adhesions has not decreased. The severe drying of the mesothelium which results from prolonged exposure of the peritoneum to dry gases (pneumoperitoneum of 2-4 hours), may give rise to a higher incidence of global peritoneal adhesions than has hitherto been encountered in open surgery. Many gynaecologists with long experience of laparoscopic surgery consider that both open and closed surgery have equally high incidences of adhesions.

20

WO 92/21354 describes a surgical adhesion as the attachment of organs or tissues to each other through scar tissue. A formation of scar tissue is described as a normal sequel to surgery or other tissue injury and is required for proper wound healing. In some cases, however, the scar tissue overgrows the intended region and creates surgical adhesions. These scar tissue surgical adhesions restrict the normal mobility and function of affected body parts. The invention disclosed in WO 92/21354 is based on the discovery that anionic polymers effectively inhibit invasion of cells associated with detrimental healing processes, ie, fibrosis, and scarring. In particular, certain inhibitory anionic polymers are useful to inhibit fibroblast invasion, thus regulating the healing process and preventing fibrosis. Anionic polymers specified in WO 92/21354 include dextran sulfate, pentosan polysulfate as well as natural

22-06-2000

PCT/GB99/01306

DESC

proteoglycans, or the glycosaminoglycan moieties of proteoglycans, including dermatan sulfate, chondroitin sulfate, keratan sulfate, heparan-sulfate, heparin and alginate.

5 By attempting to inhibit fibroblast invasion, the approach of WO 92/21354 is one of post-adhesion treatment since fibroblast invasion is a later stage, that is to say, it occurs after formation of the adhesion. The invention of WO 92/21354 attempts to prevent the adhesion becoming permanent. By contrast the present invention is concerned with the prevention of the occurrence of an adhesion.

10 According to a first aspect of the present invention there is provided a method of preventing or reducing the incidence of adhesions in or associated with a body cavity comprising an aqueous formulation containing the polysaccharide dextrin in an amount effective to prevent or reduce such adhesions, wherein the dextrin contains more than 1.5% of polymers with a degree of polymerisation (DP) greater than 12 and
15 acts as an osmotic agent to maintain a volume of the aqueous formulation in the body cavity serving to separate tissues which otherwise may adhere to each other.

The term "dextrin" means a glucose polymer which is produced by the hydrolysis of starch and which consists of glucose units linked together by means mainly of α -1,4
20 linkages. Typically dextrans are produced by the hydrolysis of starch obtained from various natural products such as wheat, rice, maize and tapioca. In addition to α -1,4 linkages, there may be a proportion of α -1,6 linkages in a particular dextrin, the amount depending on the starch starting material. Since the rate of biodegradability of α -1,6 linkages is typically less than that for α -1,4 linkages, it is preferred that, for
25 many applications, the percentage of α -1,6 linkages is less than 10% and more preferably less than 5%.

Any dextrin is a mixture of polyglucose molecules of different chain lengths. As a result no single number can adequately characterise the molecular weight of such a
30 polymer. Accordingly, various averages are used, the most common being the weight average molecular weight (Mw) and the number average molecular weight (Mn). Mw is particularly sensitive to changes in the high molecular weight content

22-06-2000

PCT/GB99/01306

DESC

of a polymer whilst M_n is largely influenced by changes in the low molecular weight content of the polymer.

It is preferred that the M_n of the dextrin is in the range of from 1,000 to 30,000 and ideally the M_w is in the range of from 3,000 to 50,000. More preferably, the M_n is from 3,000 to 8,000 and the M_w is from 5,000 to 50,000.

The term "degree of polymerisation" (DP) can also be used in connection with polymer mixtures. For a single polymer molecule, DP means the number of polymer units. For a mixture of molecules of different DP's, weight average DP and number average DP correspond to M_w and M_n . In addition, DP can also be used to characterise a polymer by referring to the polymer mixture having a certain percentage of polymers of DP greater than a particular number or less than a particular number.

It is preferred that the dextrin contains more than 15% of polymers of DP greater than 12 and, more preferably, more than 50% of polymers of DP greater than 12.

The dextrin used in the present invention is water soluble or at least forms a solution in water or a gel formulation. The dextrin used in this invention may be in the form of either unsubstituted dextrin (as obtained by the hydrolysis of starch) or may be substituted by one or more different groups. The substituents may be negatively charged groups, for instance, sulfate groups, neutral groups, or positively charged groups, for instance, quaternary ammonium groups. In the case where the substituent group is sulfate, it is preferred that the sulfated polysaccharide contains at least one sulfate group per saccharide (glucose) unit.

The present invention also provides a composition for preventing or reducing the incidence of adhesions in or associated with a body cavity comprising an aqueous formulation containing the polysaccharide dextrin in an amount effective to prevent or reduce such adhesions, wherein the dextrin contains more than 15% of polymers

22-06-2000

PCT/GB99/01306

DESC

with a degree of polymerisation (DP) greater than 12 and acts as an osmotic agent to maintain a volume of the aqueous formulation in the body cavity serving to separate tissues which otherwise may adhere to each other.

- 5 The present invention further provides the use of a composition for preventing or reducing the incidence of adhesions in or associated with a body cavity comprising an aqueous formulation containing the polysaccharide dextrin in an amount effective to prevent or reduce such adhesions, wherein the dextrin contains more than 15% of polymers with a degree of polymerisation (DP) greater than 12 and acts as an osmotic agent to maintain a volume of the aqueous formulation in the body cavity serving to separate tissues which otherwise may adhere to each other.

- 15 The present invention further provides the use of the polysaccharide dextrin in the manufacture of a composition comprising an aqueous solution or gel formulation of dextrin for preventing or reducing adhesions in humans and animals.

- 20 Dextrin is a useful material for the production of an adhesion-preventing composition because, *inter alia*, it is non-toxic, cheap and has the ability to hold fluid in a body cavity. It is also readily metabolised within the body.

- Preferably, a composition of the invention is applied to the appropriate body cavity or area after the operation has been carried out.

- 25 Preferably, the composition of the present invention is allowed to remain in the body cavity for a minimum of 2 to 3 days and especially over the period during which fibrin exudation is at a maximum. More preferably, the composition should remain in the body cavity for a period of up to 7 to 8 days in order to allow restoration of non-stick surfaces (mesothelium regeneration).

- 30 Preferably, a composition of the invention should be applied to the body cavity in a volume large enough to keep the surfaces apart. For the peritoneum, the volume should preferably be in the range 500-2000 ml and, more preferably, about 1000 ml-1500 ml.

22-06-2000

PCT/GB99/01306

DESC

Preferably, the composition should be applied to the appropriate body cavity or area in differing concentrations ideally over a concentration range of 2.5-18% and more ideally over a concentration range of 3-5% and most ideally at about 4% by weight, said concentration range is selected for a specified time span, even more ideally the concentration range is selectively altered over a period of time.

Preferably, the composition should include a concentration of dextrin which is such that the fluid largely holds in place over the period it resides in the cavity. Where a composition includes 4% by weight of dextrin then a suitable dwell period for one infusion might be of the order of 2 to 3 days. A high concentration is liable to cause ingress of fluid. A second infusion at day 3 may extend the total dwell period from 6 to 7 days.

Alternatively, a composition having a dextrin concentration of from 12 to 15% by weight may be used in a smaller volume (perhaps about 750 ml) and will be subject to ingress of fluid. However a single infusion might be sufficient for the full 6 to 7 day period.

Comparing dextrin with dextran, the latter has relatively poor biocompatibility. It is subject to immunological hypersensitivity due to its concentration in lymph nodes and its lack of metabolisability. At best, a dextran solution or suspension will act not so much to separate surfaces and therefore prevent adhesions but simply as a lubricant. Dextrin advantageously serves as an osmotic agent, which can maintain the volume of a solution in the peritoneal cavity. The continued presence of the dextrin solution within the cavity serves to separate tissues which otherwise may adhere to each other.

The use of a solution or gel formulation of dextrin is also advantageous by comparison with a prior art technique which makes use of synthetic films in the form of patches which are applied to particular areas where maximum damage has occurred. However, in the case of a body cavity, such as the peritoneum, the damage is liable to occur as well at a distance from the operative site, especially in

22-06-2000

PCT/GB99/01306

DESC

laparoscopy, due to the drying which takes place. In some instances global damage over an area of as much as two square metres can take place.

In responding to a wound, the body causes circulating fibrinogen to form fibrin and it

5 is this production of fibrin which is associated with the formation of adhesions. Calcium ions are required to polymerise fibrinogen to fibrin and, accordingly, a composition of the present invention may include a calcium binding agent such as EDTA or sodium citrate.

10 A composition of the present invention may include a suitable lubricant such as a phosphospholipid.

A composition of the present invention may include a hyaluronate or
glycocosolaminoglycan or a material which is associated with serosal lubrication and
15 which has strong anti-adhesive properties. In this case the dextrin solution or gel formulation is effective in spreading the hyaluronate throughout the whole peritoneum.

A composition of the present invention may include an antibiotic agent or a
20 material/agent which is associated with preventing an infection, or build up of bacteria or foreign bodies or the like. A composition including such a material/agent would be particularly advantageous in prevention or amelioration of pelvic inflammatory disease.

25 A composition of the present invention may also include a fibrinolytic agent or an analogue thereof, an anti-inflammatory agent or an analogue thereof, dextrin sulphate and/or methylene blue.

The present invention provides a preferred composition comprising an aqueous
30 solution or gel formulation of dextrin, one or more phosphospholipids and

22-06-2000

PCT/GB99/01306

DESC

hyaluronate. Such a composition is not only highly effective in preventing adhesions but also has a good shelf life.

- 5 Mesothelial secretion of prostacyclin has been demonstrated and this activity enhances the non-stick properties of the mesothelium. The present invention provides a composition comprising dextrin together with prostacyclin or an analogue thereof.

- 10 According to a further aspect of the invention there is provided a biocompatible, bioresorbable, and non-toxic adhesion prevention kit for surgical use in humans or animals, comprising an aqueous solution or gel formulation of dextrin as hereinbefore described, and optionally or additionally comprising a calcium binding agent as hereinbefore described and/or a suitable lubricant as hereinbefore described and/or prostacyclin or an analogue thereof as hereinbefore described and/or an antibiotic agent as hereinbefore described..

15

EVIDENCE IN SUPPORT OF THE INVENTION

PROTOCOL:

- 20 Animals: One hundred thirty, female New Zealand White rabbits, 2.4-2.7 kg, were purchased from Irish Farms (Norco, CA) and quarantined in the USC Vivaria for at least 2 days prior to use. Ten rabbits were randomised into thirteen treatment groups prior to initiation of surgery. The rabbits were housed on a 12:12 light:dark cycle with food and water available *ad libitum*.

- 25 Materials: The solutions (7.5% [wt/vol] icodextrin-Lot # 98A06G33, 20% [wt/vol] icodextrin-Batch # SP184772 and placebo (electrolyte solution for icodextrin)-Batch # SP184829 were supplied by ML Laboratories Plc. Icodextrin is a [1 → 4] - α - Glucan having more than 85% of its molecules with molecular weights between 1,640 - 45,000 with a weight average molecular weight of approximately 20,000. The placebo electrolyte
30 solution contained 5.4g sodium chloride, 4.5g sodium lactate, 257 mg calcium chloride, 51 mg magnesium chloride in 1 litre water for injection. The sutures used to close the muscle and skin were 3-0 coated Dexon II suture (Davis and Geck, Manati, PR).

Double Uterine Horn Model: Rabbits were anaesthetised with a mixture of 55 mg/kg ketamine hydrochloride and 5 mg/kg Rompum intramuscularly. Following preparation for sterile surgery, a midline laparotomy was performed. The uterine horns were exteriorised and traumatised by abrasion of the serosal surface with gauze until punctuate bleeding developed. Ischaemia of both uterine horns was induced by removal of the collateral blood supply. The remaining blood supply to the uterine horns was the ascending branches of the utero-vaginal arterial supply of the myometrium. At the end of surgery, 10 to 75 ml (10, 25, 50, 75 ml) of 7.5% or 20% icodextrin, 10 or 75 ml placebo or no treatment (control) was administered. After 7 days, the rabbits were terminated and the percentage of the area of the horns adherent to the various organs was determined. In addition, the tenacity of the adhesions was scored using the following system:

- 0 = No adhesions;
- 1 = Mild, easily dissectable adhesions;
- 2 = Moderate adhesions; non-dissectable, does not tear the organ;
- 3 = Dense adhesions; non-dissectable, tears organ when removed.

In addition an overall score which takes into account all of the above data was given to each rabbit. The following scoring system was used:

- 0 No adhesions;
- 0.5+ Light, filmy pelvic adhesions involving only one organ, typically only 1 or 2 small adhesions;
- 1.0+ Light, filmy adhesions, not extensive although slightly more extensive than 0.5;
- 1.5+ Adhesions slightly tougher and more extensive than a 1 rating;
- 2.0+ Tougher adhesions, a little more extensive, uterine horns usually have adhesions to both bowel and bladder;
- 2.5+ Same as 2, except the adhesions are usually not filmy at any site and more extensive;

- 3.0+ Tougher adhesions than 2, more extensive, both horns are attached to the bowel and bladder, some movement of the uterus possible;
- 3.5+ Same as 3, but adhesions slightly more extensive and tougher;
- 4.0+ Severe adhesions, both horns attached to the bowel and bladder, unable to move the uterus without tearing the adhesions.

The rabbits were scored by two independent observers that were blinded to the prior treatment of the animal. If there was disagreement as to the score to be assigned to an individual animal, the higher score was given.

Statistical Analysis: The tenacity and overall scores were analysed by rank order analysis and analysis of variance on the ranks. The percentage area of the horns involved to the various organs was compared by Student's t test. The data from the incidence of adhesion formation was analysed by Chi square analysis. The comparison with placebo shown on Table 14 was done between the 10 ml placebo group and data from animals which received 10-25 ml of icodextrin or between the 75 ml placebo group and data from animals which received 50 or 75 ml icodextrin.

RESULTS: One rabbit from the group treated with 50 ml 20% icodextrin died postoperatively without evidence of inflammation or oedema at necropsy and was replaced. During the postoperative evaluation of the rabbits, it was noted that several rabbits given the higher volumes of icodextrin had "bulging" abdomens for the first few postoperative days. This occurred in 3 rabbits which received 75 ml or 7.5% icodextrin and 8 rabbits which received 75 ml of 20% icodextrin. The bulging was observed for 24 hours in the rabbits which received 7.5% icodextrin and 48-72 hours in the rabbits which received 20% icodextrin. This bulging was not observed in the group of rabbits which received 75 ml of placebo. No excess fluid was observed in any icodextrin or placebo-treated rabbits at necropsy. One rabbit, which received 75 ml of 20% icodextrin, had a small amount of subcutaneous fluid at necropsy.

The effect of icodextrin on the formation of adhesions in this rabbit model can be found in Tables 1-13. The effect of icodextrin on the incidence of adhesions can be found in Table 14. For each site, the extent and tenacity (tenacity in parentheses) of the adhesions between the horn and that site were given. In the final row of each column (with the exception of the column on the far right), the mean and standard error of the mean for the extent score for each site is given. In the final row of the final column, the mean and standard error of the mean of the ranks is given. If an extent or rank order was reduced compared to control ($p \leq 0.05$), a * is in the appropriate row. At higher volumes (25 to 75 ml) of icodextrin, there was a significant reduction in the formation of adhesions. However, no difference between the 7.5% and 20% solutions was noted in this study. This efficacy is in the absence of inflammation noted with some materials implanted intraperitoneally.

In conclusion results demonstrated that high volumes of icodextrin (both percentages) were highly efficacious in the reduction of adhesion formation in this model with efficacy noted after administration of 50 ml or 75 ml of icodextrin. The lower volumes of icodextrin have less effect and the placebo had no effect on adhesion formation. Thus we have demonstrated that the composition of the present invention is effective in reducing the incidence of post-operative adhesion formation.

Table 1. Data from Surgical control Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
30(2)	30(2)	30(1)	40(1)	30(2)	30(2)	30(1)	40(1)	2.5+
30(1)	30(1)	50(2)	50(2)	30(1)	30(1)	30(1)	50(2)	2.5+
30(2)	30(1)	40(2)	40(2)	30(2)	30(2)	40(2)	40(2)	3.0+
40(1)	20(1)	50(2)	30(1)	40(1)	20(1)	30(1)	30(1)	3.0+
20(1)	30(1)	50(2)	40(2)	20(1)	30(1)	50(2)	40(2)	3.0+
40(1)	30(1)	50(1)	40(1)	40(1)	30(1)	60(1)	40(1)	3.5+
40(1)	-	50(1)	40(2)	40(1)	-	50(1)	40(1)	3.0+
40(1)	20(1)	50(1)	40(1)	40(1)	20(1)	40(2)	40(1)	3.0+
40(2)	20(2)	40(2)	30(2)	40(2)	20(2)	50(1)	30(2)	3.5+
40(1)	20(1)	60(1)	50(1)	40(1)	20(1)	60(1)	50(1)	3.0+
31±3.7	23±3.0	47±2.6	40±2.1	34±3.7	23±3.0	44±3.7	40±2.1	111.2±4.0

Table 2. Data from 10ml Placebo Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
30(1)	20(1)	50(2)	30(1)	30(1)	20(1)	40(2)	30(1)	2.5+
50(2)	50(2)	60(1)	30(2)	50(2)	50(2)	60(1)	30(2)	3.5+
40(2)	-	50(1)	20(2)	40(2)	-	50(1)	20(2)	3.0+
20(1)	-	30(1)	20(1)	20(1)	-	30(1)	20(2)	1.5+
-	30(2)	40(1)	40(2)	-	30(2)	40(1)	40(2)	2.5+
50(1)	20(1)	40(2)	30(1)	50(1)	20(1)	50(1)	30(1)	3.0+
30(2)	-	20(2)	40(2)	30(2)	-	20(2)	40(2)	3.0+
30(2)	-	30(2)	40(1)	30(2)	-	50(2)	40(1)	3.0+
50(2)	-	40(1)	50(2)	50(2)	-	40(1)	50(2)	3.0+
30(2)	20(2)	30(1)	40(1)	30(2)	20(2)	50(1)	40(1)	2.5+
33±5.0	14±5.4	39±3.9	34±3.0	33±5.0	14±5.4	43±3.7	34±3.1	100.7±7.5

Table 3. Data from 75ml Placebo Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
40(2)	30(2)	50(2)	40(2)	40(20)	30(2)	40(3)	40(2)	3.5+
-	-	50(1)	20(1)	-	-	40(1)	20(1)	2.5+
-	40(2)	30(10)	40(2)	-	40(2)	40(2)	40(2)	3.0+
40(1)	20(1)	50(1)	20(1)	40(1)	20(1)	50(1)	20(1)	3.0+
20(1)	-	40(1)	20(1)	20(1)	-	30(1)	20(1)	2.0+
-	10(1)	20(1)	40(1)	-	10(1)	40(1)	40(1)	2.0+
-	30.2	50(2)	40(2)	-	30(2)	50(2)	40(2)	3.0+
40(1)	20(1)	50(1)	20(1)	40(1)	20(1)	30(1)	20(1)	2.5+
20(1)	-	60(1)	50(1)	20(1)	-	50(1)	50(1)	3.0+
20(2)	10(1)	40(1)	30(1)	20(2)	10(1)	50(2)	30(1)	3.0+
18±5.5	16±4.5	44±3.7	32±3.6	18±5.5	16±4.5	42±7.9	32±3.6	100.5±6.8

Table 4. Data from 10ml 7.5% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
30(1)	20(1)	50(1)	40(1)	30(1)	20(1)	50(1)	40(1)	2.5+
40(1)	30(1)	50(1)	40(2)	40(1)	30(1)	50(1)	40(2)	3.0+
40(1)	10(1)	30(1)	10(1)	40(1)	10(1)	10(1)	10(1)	2.0+
30(1)	20(1)	30(2)	30(1)	30(1)	20(1)	30(2)	30(1)	2.5+
-	-	10(1)	10(1)	-	-	10(1)	10(1)	1.0+
40(2)	20(1)	50(1)	30(1)	40(2)	20(1)	50(2)	30(1)	3.5+
-	10(1)	40(1)	40(2)	-	10(1)	50(1)	40(2)	2.5+
-	30(2)	30(1)	40(2)	-	30(2)	50(2)	40(2)	3.0+
30(2)	-	50(1)	30(1)	30(2)	-	40(1)	30(1)	2.5+
30(2)	-	10(1)	30(1)	30(2)	-	50(1)	30(1)	2.5+
24±5.4	14±3.7	35±5.0	30±3.7	24±5.4	14±3.7	39±5.3	30±3.7	88.8±9.3 *

Table 5. Data from 15ml 7.5% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
20(1)	10(1)	40(1)	40(1)	20(1)	10(1)	40(1)	40(1)	2.5+
10(1)	-	30(1)	30(1)	10(1)	-	30(1)	30(1)	2.0+
30(2)	30(2)	40(1)	20(1)	30(2)	30(2)	40(1)	20(2)	2.5+
10(1)	20(1)	30(1)	10(1)	10(1)	20(1)	30(1)	10(1)	2.0+
30(1)	30(1)	40(1)	30(1)	30(1)	30(1)	40(1)	30(1)	2.5+
40(1)	10(1)	50(1)	50(1)	40(1)	10(1)	50(1)	50(1)	3.0+
-	20(1)	30(1)	20(1)	-	20(1)	30(1)	20(1)	1.5+
20(1)	10(1)	30(1)	10(1)	20(1)	10(1)	30(1)	10(1)	1.5+
30(2)	30(2)	40(1)	10(1)	30(2)	30(2)	50(1)	10(1)	2.5+
-	30(1)	40(1)	30(2)	-	30(1)	50(2)	30(2)	2.5+
19±4.3 *	19±3.5	37±2.1 *	25±4.3 *	19±4.3 *	19±3.5	39±2.8 *	25±4.3 *	78.2±7.0 *

Table 6. Data from 25ml 7.5% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
10(1)	-	30(1)	20(1)	10(1)	-	30(1)	20(1)	1.5+
-	-	40(1)	-	-	-	40(1)	-	1.0+
10(1)	-	30(1)	20(2)	10(1)	-	30(1)	20(1)	2.0+
10(1)	-	30(1)	10(1)	10(1)	-	30(1)	10(1)	2.0+
-	-	10(1)	10(1)	-	-	-	10(1)	1.0+
-	-	30(1)	-	-	-	10(1)	-	1.0+
-	-	20(1)	40(1)	-	-	30(2)	40(1)	2.0+
40(1)	30(1)	30(1)	10(1)	40(1)	30(1)	30(1)	10(1)	2.5+
10(1)	-	20(1)	10(1)	10(1)	-	30(1)	10(1)	1.5+
30(1)	-	30(1)	30(1)	30(1)	-	40(1)	30(1)	2.0+
11±4.3 *	3±3.0 *	27±2.6 *	15±4.0 *	11±4.3 *	3±3.0 *	27±3.0 *	15±4.0 *	50.6±7.6 *

Table 7. Data from 50ml 7.5% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
-	-	30(1)	10(1)	-	-	30(1)	10(1)	1.0+
-	-	20(1)	10(1)	-	-	20(1)	10(1)	1.0+
10(1)	-	30(1)	-	10(1)	-	30(1)	-	1.0+
-	-	20(1)	10(1)	-	-	20(1)	10(1)	1.0+
-	-	30(2)	10(1)	-	-	30(2)	10(1)	1.5+
20(1)	10(1)	10(1)	-	20(1)	-	-	-	1.0+
-	10(1)	30(2)	40(2)	-	10(1)	30(2)	40(2)	2.5+
30(1)	-	40(1)	10(1)	30(1)	-	10(1)	10(1)	2.0+
10(1)	10(2)	10(1)	10(1)	10(1)	10(2)	10(1)	10(1)	1.5+
-	-	20(1)	30(1)	-	-	40(1)	30(1)	1.5+
7±3.4 *	3±1.5 *	24±3.1 *	13±4.0 *	7±3.4 *	2±1.3 *	22±3.9 *	13±4.0 *	39.4±7.4 *

Table 8. Data from 75ml 7.5% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
-	-	-	-	-	-	10(1)	-	0.5+
10(2)	-	30(1)	-	10(2)	-	20(1)	-	1.0+
-	-	-	-	-	10(1)	30(1)	-	0.5+
10(1)	10(1)	40(1)	-	10(1)	10(1)	-	-	1.5+
20(2)	-	30(1)	-	20(2)	-	-	-	1.0+
-	-	10(1)	-	-	-	50(1)	-	1.0+
10(1)	20(1)	-	10(1)	10(1)	20(1)	10(1)	10(1)	1.5+
-	-	20(1)	10(1)	-	-	30(1)	10(1)	1.0+
-	10(1)	10(1)	-	-	10(1)	10(1)	-	1.0+
-	-	20(1)	20(1)	-	-	20(1)	20(1)	1.0+
5±2.2 *	4±2.2 *	16±4.5 *	4±2.2 *	5±2.2 *	5±2.2 *	18±4.9 *	4±2.2 *	22.5±4.2 *

Table 9. Data from 10ml 20% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
20(2)	-	50(2)	30(2)	20(2)	-	50(2)	30(2)	3.0+
10(1)	-	10(1)	10(1)	10(1)	-	50(1)	10(1)	2.0+
40(1)	-	50(2)	30(1)	40(1)	-	40(1)	30(1)	2.5+
40(1)	20(1)	40(1)	30(2)	40(1)	20(1)	30(1)	30(2)	2.5+
-	20(1)	10(1)	-	-	20(1)	20(1)	-	1.0+
30(2)	-	40(1)	30(1)	30(2)	-	30(1)	30(1)	2.5+
30(1)	20(2)	40(1)	40(1)	30(1)	20(2)	10(1)	40(1)	2.5+
10(1)	-	50(1)	20(1)	10(1)	-	30(1)	20(1)	2.0+
20(1)	10(1)	40(1)	40(1)	20(1)	10(1)	30(1)	40(1)	2.5+
20(1)	-	20(1)	20(1)	20(1)	-	40(2)	20(1)	2.0+
22±4.2	7±3.0 *	35±5.0 *	25±4.0 *	22±4.2	7±3.0 *	33±4.0	25±4.0 *	78±7.9 *

Table 10. Data from 15ml 20% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
-	-	50(1)	50(1)	-	-	50(1)	50(1)	2.5+
20(1)	20(1)	20(1)	20(1)	20(1)	20(1)	20(1)	20(1)	2.0+
-	30(2)	50(1)	30(1)	-	30(2)	20(1)	30(1)	2.5+
20(1)	20(1)	40(1)	30(1)	20(1)	20(1)	40(1)	30(1)	2.0+
30(1)	20(1)	40(1)	20(1)	30(1)	20(1)	40(1)	20(1)	2.0+
40(2)	30(2)	50(1)	50(1)	40(2)	30(2)	50(1)	50(1)	3.0+
20(1)	-	20(1)	-	-	-	-	-	0.5+
-	-	-	10(1)	-	-	-	10(1)	0.5+
-	20(1)	10(1)	10(1)	-	20(1)	20(1)	10(1)	1.5+
30(1)	30(1)	40(1)	40(1)	20(1)	30(1)	40(1)	30(1)	2.5+
16±4.8 *	17±4.0	32±5.7	25±5.2 *	14±5.0 *	17±4.0	28±5.9	25±5.2 *	61.3±11.6 *

6/7

Table 11. Data from 25ml 20% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
20(1)	-	-	10(1)	20(1)	-	40(2)	10(1)	1.5+
20(1)	20(1)	40(1)	-	-	-	30(1)	-	1.5+
10(1)	-	40(1)	10(1)	10(1)	-	30(1)	10(1)	1.5+
30(1)	-	40(1)	-	30(1)	-	10(1)	-	1.5+
-	-	10(1)	10(1)	-	-	10(1)	10(1)	1.0+
10(1)	20(1)	10(1)	10(1)	10(1)	20(1)	20(1)	10(1)	2.0+
20(2)	-	30(1)	40(2)	20(2)	-	30(1)	40(2)	2.5+
-	-	20(1)	20(1)	-	-	50(1)	20(1)	1.5+
-	-	-	-	-	-	10(1)	-	0.5+
-	20(1)	10(1)	-	-	20(1)	20(1)	-	1.0+
11±3.5 *	6±3.1 *	20±5.2	10±3.0 *	9±3.5 *	4±2.7 *	25±4.3 *	10±3.9 *	42.5±7.5 *

Table 12. Data from 50ml 20% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
-	-	10(1)	-	-	-	-	-	0.5+
10(1)	-	10(1)	-	-	-	20(2)	-	1.0+
20(1)	-	20(1)	10(1)	20(1)	-	10(1)	10(1)	1.5+
-	10(1)	-	-	-	-	-	-	0.5+
-	-	10(1)	10(1)	-	-	20(1)	10(1)	1.0+
30(1)	-	20(1)	10(1)	30(1)	-	20(1)	10(1)	1.5+
-	10(1)	-	10(1)	-	10(1)	20(1)	10(1)	1.5+
-	20(1)	40(1)	30(1)	-	20(1)	10(1)	30(1)	2.0+
-	30(1)	40(1)	10(1)	-	30(1)	20(1)	10(1)	2.0+
-	30(2)	30(1)	20(1)	-	30(2)	30(2)	20(1)	2.0+
6±3.4 *	10±3.9 *	18±4.7 *	10±3.0	5±3.4 *	9±4.1 *	15±3.1 *	10±3.0 *	38.4±7.6 *

Table 13. Data from 75ml 20% Icodextrin Rabbits
% Horn Involved

Right Horn				Left Horn				Overall
Bowel	Bladder	Itself	Left	Bowel	Bladder	Itself	Right	
-	-	10(1)	10(1)	-	-	20(2)	10(1)	1.0+
-	-	-	-	-	-	30(1)	-	0.5+
-	-	30(1)	10(1)	-	-	30(1)	10(1)	1.0+
20(1)	40(1)	10(1)	-	20(1)	40(1)	10(1)	-	2.0+
-	10(1)	30(1)	10(1)	-	10(1)	30(1)	10(1)	1.5+
10(1)	10(1)	20(1)	30(1)	10(1)	10(1)	20(1)	30(1)	1.5+
-	30(1)	10(1)	10(1)	-	30(1)	10(1)	10(1)	1.5+
-	-	40(1)	10(1)	-	-	20(1)	10(1)	1.5+
-	-	10(1)	20(1)	-	-	-	20(1)	1.0+
-	-	10(1)	-	20(1)	-	-	-	0.5+
3±2.1 *	9±4.5 *	17±4.0 *	10±3.0 *	5±2.7 *	9±4.6 *	17±3.7 *	10±3.0 *	31.7±6.3 *

Table 14. Incidence of Adhesion Formation

	# Sites Free/ # Possible	% Adhesion Free	p Value	
			Control	Placebo
Control	2/80	2.5		
10ml Placebo	12/80	15.0	0.012	
75ml Placebo	14/80	17.5	0.004	
10ml 7.5% Icodextrin	12/80	15.0	0.012	1.00
15ml 7.5% Icodextrin	6/80	7.5	0.277	0.211
25ml 7.5% Icodextrin	31/80	38.8	0.000	0.001
50ml 7.5% Icodextrin	32/80	40.0	0.000	0.003
75ml 7.5% Icodextrin	44/80	55.0	0.000	0.000
10ml 20% Icodextrin	16/80	20.0	0.001	0.533
15ml 20% Icodextrin	20/80	25.0	0.000	0.167
25ml 20% Icodextrin	34/80	42.5	0.000	0.000
50ml 20% Icodextrin	36/80	45.0	0.000	0.000
75ml 20% Icodextrin	36/80	45.0	0.000	0.000